Approximation by Modified Jain-Baskakov Operators

Vishnu Narayan Mishra[a,b,1], Preeti Sharma[a], Marius Birou[c]

[a]*Department of Applied Mathematics & Humanities, Sardar Vallabhbhai National Institute of Technology, Ichchhanath Mahadev Dumas Road, Surat -395 007 (Gujarat), India*
[b]*L. 1627 Awadh Puri Colony Beniganj, Phase -III, Opposite - Industrial Training Institute (I.T.I.), Ayodhya Main Road Faizabad-224 001, (Uttar Pradesh)*
[c]*Mathematics Department, Technical University of Cluj Napoca, Cluj Napoca, Romania*

Dedicated to the 66-th Happy Anniversary of our great Master and friend academician Prof. Dr. Heiner Gonska

arXiv:1508.05309v2 [math.FA] 9 Sep 2015

Abstract

In the present paper we discuss the approximation properties of Jain-Baskakov operators with parameter c. The present paper deals with the modified forms of the Baskakov basis functions. We establish some direct results, which include the asymptotic formula and error estimation in terms of the modulus of continuity and weighted approximation.

Keywords: Jain operators, Baskakov operators, King-type operators, Weighted approximation.
2000 MSC: 41A25, 41A36.

---

1. Introduction

In 1972, G.C. Jain [7] introduced and studied the following class of positive linear operators

$$P_n^{[\beta]}(f,x)=\sum_{v=0}^{\infty}\omega_\beta(v,nx)f\left(\frac{v}{n}\right), \tag{1.1}$$

where

$$\omega_\beta(v,nx)=nx(nx+v\beta)^{v-1}\frac{e^{-(nx+v\beta)}}{v!}, \tag{1.2}$$

and $\beta\in[0,1)$, $f\in C(\mathbb{R}_+)$. In the particular case $\beta=0$, P_n^0, $n\in\mathbb{N}$, turns into well-known Szasz-Mirakjan operators [12]. A Kantorovich-type extension of $P_n^{[\beta]}$ was given in [14]. In the last few decades several authors contributed to the approximation theory, fractional differential and integral equations have recently been applied in various areas of engineering, mathematics, physics and bio-engineering and other applied sciences (see [2], [6], [13], [15], [16], [17], [18] and references therein).

Very recently Patel and Mishra [10] introduced modified Jain-Baskakov operators as follows

$$\mathcal{D}_{n,c}^{\beta}(f,x) \quad = \quad \frac{(n-c)}{c}\sum_{v=1}^{\infty}\omega_\beta(v,nx)\int_0^\infty p_{n,v-1,c}(t)f(t)dt+e^{-nx}f(0), \tag{1.3}$$

Email addresses: `vishnunarayanmishra@gmail.com` (Vishnu Narayan Mishra), `preeti.iitan@gmail.com` (Preeti Sharma), `marius.birou@math.utcluj.ro` (Marius Birou)

[1] Corresponding author

where $p_{n,v-1,c}(t)=c\frac{\Gamma(n/c+v-1)}{\Gamma(v)\Gamma(n/c)}\cdot\frac{(ct)^{v-1}}{(1+ct)^{n/c+v-1}}$ and $\omega_\beta(v,nx)$ are as defined in (1.2).
As a special case, i.e., $c=1$, the operators (1.3) reduce to the Jain-Baskakov operators which is defined in [10]. The operators given by (1.3) have different approximation properties than the operators defined for $c=1$.

The present paper deals with the study of some direct results in ordinary approximation for the modified Jain-Baskakov operators with parameter c. We study weighted approximation theorems, rate of convergence of these operators.

2. Auxiliary Results

Lemma 1. *([7], [3]) For $P_n^{[\beta]}(t^m,x)$, $m=0,...,4$, we have*

$$P_n^{[\beta]}(1,x)=1,\quad P_n^{[\beta]}(t,x)=\frac{x}{1-\beta},\quad P_n^{[\beta]}(t^2,x)=\frac{x^2}{(1-\beta)^2}+\frac{x}{n(1-\beta)^3},$$

$$P_n^{[\beta]}(t^3,x)=\frac{x^3}{(1-\beta)^3}+\frac{3x^2}{n(1-\beta)^4}-\frac{(6\beta^4-6\beta^3-2\beta-1)x}{n^2(1-\beta)^5},$$

$$\begin{aligned}P_n^{[\beta]}(t^4,x) &= \frac{x^4}{(1-\beta)^4}+\frac{6x^3}{n(1-\beta)^5}-\frac{(36\beta^4-72\beta^3+36\beta^2-8\beta-7)x^2}{n^2(1-\beta)^6}\\ &\quad+\frac{(105\beta^5-14\beta^4-2\beta^3+12\beta^2+8\beta+1)x}{n^3(1-\beta)^7}.\end{aligned}$$

Lemma 2. *For $n>5c>0$, we have*

$$\mathcal{D}_{n,c}^{\beta}(1,x)=1,\quad \mathcal{D}_{n,c}^{\beta}(t,x)=\frac{nx}{(n-2c)(1-\beta)},$$

$$\mathcal{D}_{n,c}^{\beta}(t^2,x)=\frac{n^2}{(n-2c)(n-3c)}\left[\frac{x^2}{(1-\beta)^2}+\frac{x\left(2-2\beta+\beta^2\right)}{n(1-\beta)^3}\right],$$

$$\mathcal{D}_{n,c}^{\beta}(t^3,x)=\frac{n^2x^2(-(1-\beta)xn+3(\beta^2-2\beta+2))}{(1-\beta)^4(n-2c)(n-3c)(n-4c)}+o\left(\frac{1}{n}\right),$$

$$\mathcal{D}_{n,c}^{\beta}(t^4,x)=\frac{n^3x^3((1-\beta)xn+6(\beta^2-2\beta+2))}{(1-\beta)^5(n-2c)(n-3c)(n-4c)(n-5c)}+o\left(\frac{1}{n}\right).$$

Proof. For $n>(j+1)c$, we have

$$\begin{aligned}\int_0^\infty t^j p_{n,v-1,c}(t)dt &= \frac{c\Gamma(n/c+v-1)}{\Gamma(v)\Gamma(n/c)}\cdot\frac{\Gamma(v+j)\Gamma(n/c-j-1)}{c^{j+1}\Gamma(n/c+v-1)}\\ &= c\frac{(v+j-1)...v}{(n-c)...(n-(j+1)c)}.\end{aligned}$$

It follows that

$$\mathcal{D}_{n,c}^{\beta}(1,x)=\frac{n-c}{c}\sum_{n=1}^{\infty}\omega_\beta(v,nx)\cdot\frac{c}{n-c}+e^{-nx}=P_n^{[\beta]}(1,x),$$

$$\begin{aligned}\mathcal{D}_{n,c}^{\beta}(t,x) &= \frac{n-c}{c}\sum_{n=1}^{\infty}\omega_\beta(v,nx)\cdot\frac{cv}{(n-c)(n-2c)}\\ &= \frac{n}{n-2c}\sum_{n=1}^{\infty}\omega_\beta(v,nx)\frac{v}{n}=\frac{n}{n-2c}P_n^{[\beta]}(t,x),\end{aligned}$$

$$\begin{aligned}
\mathcal{D}_{n,c}^{\beta}(t^2,x) &= \frac{n-c}{c}\sum_{n=1}^{\infty}\omega_\beta(v,nx)\cdot\frac{cv(v+1)}{(n-c)(n-2c)(n-3c)}\\
&= \frac{n^2}{(n-2c)(n-3c)}\sum_{n=1}^{\infty}\omega_\beta(v,nx)\frac{v^2+v}{n^2}\\
&= \frac{n^2}{(n-2c)(n-3c)}\left(P_n^{[\beta]}(t^2,x)+\frac{1}{n}P_n^{[\beta]}(t,x)\right),
\end{aligned}$$

$$\begin{aligned}
\mathcal{D}_{n,c}^{\beta}(t^3,x) &= \frac{n-c}{c}\sum_{n=1}^{\infty}\omega_\beta(v,nx)\cdot\frac{cv(v+1)(v+2)}{(n-c)(n-2c)(n-3c)(n-4c)}\\
&= \frac{n^3}{(n-2c)(n-3c)(n-4c)}\sum_{n=1}^{\infty}\omega_\beta(v,nx)\frac{v^3+3v^2+2v}{n^3}\\
&= \frac{n^3}{(n-2c)(n-3c)(n-4c)}\left(P_n^{[\beta]}(t^3,x)+\frac{3}{n}P_n^{[\beta]}(t^2,x)+\frac{2}{n^2}P_n^{[\beta]}(t,x)\right),
\end{aligned}$$

$$\begin{aligned}
\mathcal{D}_{n,c}^{\beta}(t^4,x) &= \frac{n-c}{c}\sum_{n=1}^{\infty}\omega_\beta(v,nx)\cdot\frac{cv(v+1)(v+2)(v+3)}{(n-c)(n-2c)(n-3c)(n-4c)(n-5c)}\\
&= \frac{n^4}{(n-2c)(n-3c)(n-4c)(n-5c)}\sum_{n=1}^{\infty}\omega_\beta(v,nx)\frac{v^4+6v^3+11v^2+6v}{n^4}\\
&= \frac{n^4}{(n-2c)(n-3c)(n-4c)(n-5c)}\times\\
&\quad\times\left(P_n^{[\beta]}(t^4,x)+\frac{6}{n}P_n^{[\beta]}(t^3,x)+\frac{11}{n^2}P_n^{[\beta]}(t^2,x)+\frac{6}{n^3}P_n^{[\beta]}(t,x)\right).
\end{aligned}$$

Using Lemma 1 we get the conclusion. □

Lemma 3. *For $n > 5c > 0$, we have*

$$\begin{aligned}
\mu_{1,n,c}^{\beta} &= \mathcal{D}_{n,c}^{\beta}(t-x,x)=\frac{x(n\beta+2c(1-\beta))}{(n-2c)(1-\beta)},\\
\mu_{2,n,c}^{\beta} &= \mathcal{D}_{n,c}^{\beta}((t-x)^2,x)\\
&= \frac{x^2\left(n^2\beta^2+n\left(c+4c\beta-5c\beta^2\right)+6c^2-12c^2\beta+6c^2\beta^2\right)}{(n-2c)(n-3c)(1-\beta)^2}+\frac{nx\left(2-2\beta+\beta^2\right)}{(n-2c)(n-3c)(1-\beta)^3},\\
\mu_{4,n,c}^{\beta} &= \mathcal{D}_{n,c}^{\beta}((t-x)^4,x)\\
&= \frac{(1-\beta)n^3\beta^2x^4(2c(3+4\beta-7\beta^2)+\beta^2n)+6n^3\beta^2x^2(\beta^2-2\beta+2)}{(n-2c)(n-3c)(n-4c)(n-5c)(1-\beta)^5}+o\left(\frac{1}{n}\right).
\end{aligned}$$

Proof. The proof follows from Lemma 2. □

3. Direct theorem

Let the space $C_B[0,\infty)$ of all continuous and bounded functions be endowed with the norm $\|f\| = \sup\{|f(x)| : x\in[0,\infty)\}$. Further let us consider the following K-functional:

$$\mathcal{K}_2(f,\delta)=\inf_{g\in W^2}\{\|f-g\|+\delta\|g''\|\}, \tag{3.1}$$

where $\delta > 0$ and $W^2 = \{g \in C_B[0,\infty) : g', g'' \in C_B[0,\infty)\}$. By the methods as given in [11], there exists an absolute constant $M > 0$ such that

$$\mathcal{K}_2(f,\delta) \leq M\omega_2(f,\sqrt{\delta}), \tag{3.2}$$

where

$$\omega_2(f,\sqrt{\delta}) = \sup_{0<h\leq\sqrt{\delta}} \ \sup_{x\in[0,\infty)} |f(x+2h) - 2f(x+h) + f(x)|$$

is the second order modulus of smoothness of $f \in C_B[0,\infty)$.

Theorem 1. *For $f \in C_B[0,\infty)$ and $n > 3c > 0$, we have*

$$|\mathcal{D}^{\beta}_{n,c}(f,x) - f(x)| \ \leq \ \omega\left(f, \mu^{\beta}_{1,n,c}(x)\right) + M\omega_2\left(f, \sqrt{\left(\mu^{\beta}_{1,n,c}(x)\right)^2 + \mu^{\beta}_{2,n,c}(x)}\right),$$

where M is a positive constant.

Proof. We are introducing the auxiliary operators as follows

$$\hat{\mathcal{D}}^{\beta}_{n,c}(f,x) = \mathcal{D}^{\beta}_{n,c}(f,x) - f\left(x + \frac{(2c-2c\beta+n\beta)x}{(n-2c)(1-\beta)}\right) + f(x).$$

Let $g \in W^2_\infty$ and $x, t \in [0,\infty)$. By Taylor's expansion we have

$$g(t) = g(x) + (t-x)g'(x) + \int_x^t (t-u)g''(u)du.$$

Applying $\hat{\mathcal{D}}^{\beta}_{n,c}$, we get

$$\hat{\mathcal{D}}^{\beta}_{n,c}(g,x) - g(x) = g'(x)\hat{\mathcal{D}}^{\beta}_{n,c}((t-x),x) + \hat{\mathcal{D}}^{\beta}_{n,c}\left(\int_x^t (t-u)g''(u)du, x\right).$$

Now,

$$\begin{aligned}
|\hat{\mathcal{D}}^{\beta}_{n,c}(g,x) - g(x)| &\leq \hat{\mathcal{D}}^{\beta}_{n,c}\left(\left|\int_x^t (t-u)g''(u)du\right|, x\right)\\
&\leq \mathcal{D}^{\beta}_{n,c}\left((t-x)^2, x\right)\|g''\|\\
&\quad + \left|\int_x^{x+\frac{(2c-2c\beta+n\beta)x}{(n-2c)(1-\beta)}} \left(x + \frac{(2c-2c\beta+n\beta)x}{(n-2c)(1-\beta)} - u\right) g''(u)du\right|\\
&\leq \left[\mu^{\beta}_{2,n,c}(x) + \left(\frac{(2c-2c\beta+n\beta)x}{(n-2c)(1-\beta)}\right)^2\right]\|g''\|.
\end{aligned}$$

Since,

$$|\mathcal{D}^{\beta}_{n,c}(f,x)| \leq (n-c)/c \sum_{v=1}^{\infty} w_\beta(v,nx)\int_0^\infty p_{n,v-1,c}(t)|f(t)|dt + e^{-nx}|f(0)| \leq \|f\|,$$

$$\begin{aligned}
|\mathcal{D}^{\beta}_{n,c}(f,x) - f(x)| &\leq |\hat{\mathcal{D}}^{\beta}_{n,c}(f-g,x) - (f-g)(x)| + |\hat{\mathcal{D}}^{\beta}_{n,c}(g,x) - g(x)|\\
&\quad + \left|\left(x + \frac{(2c-2c\beta+n\beta)x}{(n-2c)(1-\beta)}\right) - f(x)\right|\\
&\leq 2\|f-g\| + \left[\mu^{\beta}_{2,n,c}(x) + \left(\frac{(2c-2c\beta+n\beta)x}{(n-2c)(1-\beta)}\right)^2\right]\|g''\|\\
&\quad + \omega\left(f, \frac{(2c-2c\beta+n\beta)x}{(n-2c)(1-\beta)}\right).
\end{aligned}$$

Taking infimum overall $g \in W^2$, we get

$$\begin{aligned} |\mathcal{D}_{n,c}^{\beta}(f,x)-f(x)| &\leq \mathcal{K}_2\left(f,\mu_{2,n,c}^{\beta}(x)+\left(\frac{(2c-2c\beta+n\beta)x}{(n-2c)(1-\beta)}\right)^2\right) \\ &\quad +\omega\left(f,\frac{(2c-2c\beta+n\beta)x}{(n-2c)(1-\beta)}\right). \end{aligned}$$

In view of (3.2), we have

$$\begin{aligned} |\mathcal{D}_{n,c}^{\beta}(f,x)-f(x)| &\leq M\omega_2\left(f,\sqrt{\mu_{2,n,c}^{\beta}(x)+\left(\frac{(2c-2c\beta+n\beta)x}{(n-2c)(1-\beta)}\right)^2}\right) \\ &\quad +\omega\left(f,\frac{(2c-2c\beta+n\beta)x}{(n-2c)(1-\beta)}\right), \end{aligned}$$

which proves the Theorem. □

3.1. Rate of Convergence

Let $B_{\rho_0}(\mathbb{R}^+)$ be the set of all functions f defined on $\mathbb{R}^+$ satisfying the condition $|f(x)| \leq M_f(1+x^2)$, where M_f is a constant depending only on f. Now, $C_{\rho_0}(\mathbb{R}^+)$ can be represented as $\{f \in B_{\rho_0}(\mathbb{R}^+) : f$ is continuous and $\frac{f(x)}{\rho_0(x)}$ is convergent as $x \to \infty\}$. For any positive a, by

$$\omega_a(f,\delta)=\sup_{|t-x|\leq\delta}\ \sup_{x,t\in[0,a]}|f(t)-f(x)|$$

we denote the usual modulus of continuity of f on the closed interval $[0,a]$. We know that for a function $f \in C_{\rho_0}(\mathbb{R}^+)$, the modulus of the continuity $\omega_a(f,\delta)$ tends to zero.

Now, we give a rate of convergence theorem for the Modified Jain-Baskakov operators.

Theorem 2. *Let $f \in C_{\rho_0}(\mathbb{R}^+)$ and $\omega_{a+1}(f,\delta)$ be its modulus of continuity on the finite interval $[0,a+1] \subset \mathbb{R}^+$, where $a > 0$. Then for $n > 3c$,*

$$\|\mathcal{D}_{n,c}^{\beta}(f)-f\|_{C[0,a]} \leq 6M_f(1+a^2)\mu_{2,n,c}^{\beta}(x)+2\omega_{a+1}\left(f,\sqrt{\mu_{2,n,c}^{\beta}(x)}\right).$$

Proof. For $x \in [0,a]$ and $t > a+1$, since $t-x > 1$, we have

$$\begin{aligned} |f(t)-f(x)| &\leq M_f(2+x^2+t^2) \\ &\leq M_f\left(2+3x^2+2(t-x)^2\right) \\ &\leq 6M_f(1+a^2)(t-x)^2. \qquad (3.3) \end{aligned}$$

For $x \in [0,a]$ and $t \leq a+1$, we have

$$|f(x)-f(t)| \leq \omega_{a+1}(f,|t-x|) \leq \left(1+\frac{|t-x|}{\delta}\right)\omega_{a+1}(f,\delta) \qquad (3.4)$$

with $\delta > 0$.

From (3.3) and (3.4) we can write

$$|f(t)-f(x)| \leq 6M_f(1+a^2)(t-x)^2+\left(1+\frac{|t-x|}{\delta}\right)\omega_{a+1}(f,\delta), \qquad (3.5)$$

for $x \in [0,a]$ and $t \geq 0$. Thus

$$\begin{aligned} |\mathcal{D}^{\beta}_{n,c}(f,x) - f(x)| &\leq \mathcal{D}^{\beta}_{n,c}\left(|f(t)-f(x)|, x\right) \\ &\leq 6M_f(1+a^2)\mathcal{D}^{\beta}_{n,c}((t-x)^2,x) \\ &\quad +\omega_{a+1}(f,\delta)\left(1+\frac{1}{\delta}\mathcal{D}^{\beta}_{n,c}((t-x)^2,x)^{1/2}\right). \end{aligned}$$

Hence, by Schwarz's inequality and Lemma 3, for $x \in [0,a]$

$$|\mathcal{D}^{\beta}_{n,c}(f,x) - f(x)| \leq 6M_f(1+a^2)\mu^{\beta}_{2,n,c}(x) + \omega_{a+1}(f,\delta)\left(1+\frac{1}{\delta}\sqrt{\mu^{\beta}_{2,n,c}(x)}\right).$$

By taking $\delta = \sqrt{\mu^{\beta}_{2,n,c}(x)}$, we get

$$\|\mathcal{D}^{\beta}_{n,c}(f) - f\|_{C[0,a]} \leq 6M_f(1+a^2)\mu^{\beta}_{2,n,c}(x) + 2\omega_{a+1}\left(f, \sqrt{\mu^{\beta}_{2,n,c}(x)}\right).$$

This completes the proof of Theorem. $\square$

3.2. Weighted approximation

Now we shall discuss the weighted approximation theorem, where the approximation formula holds true on the interval $\mathbb{R}^+$.

Theorem 3. *If $f \in C_{\rho_0}(\mathbb{R}^+)$, $\lim_{n\to\infty} \beta_n = 0$ and $n > 3c$. Then,*

$$\lim_{n\to\infty} \|\mathcal{D}^{\beta_n}_{n,c}(f) - f\|_{\rho_0} = 0.$$

Proof. Using the theorem in [4] we see that it is sufficient to verify the following three conditions

$$\lim_{n\to\infty} \|\mathcal{D}^{\beta_n}_{n,c}(t^r,x) - x^r\|_{\rho_0} = 0, \quad r = 0,1,2. \tag{3.6}$$

Since $\mathcal{D}^{\beta_n}_{n,c}(1,x) = 1$, the first condition of (3.6) is satisfied for $r = 0$.

Now, for $n > 2c$

$$\begin{aligned} \|\mathcal{D}^{\beta_n}_{n,c}(t,x) - x\|_{\rho_0} &= \sup_{x\in[0,\infty)} \frac{|\mathcal{D}^{\beta_n}_{n,c}(t,x) - x|}{1+x^2} \\ &\leq \left|\frac{n}{(n-2c)(1-\beta_n)} - 1\right| \sup_{x\in[0,\infty)} \frac{x}{1+x^2} \\ &\leq \frac{2c}{(n-2c)} + \frac{n\beta_n}{(n-2c)(1-\beta_n)} \end{aligned}$$

which implies $\|\mathcal{D}^{\beta_n}_{n,c}(t,x) - x\|_{\rho_0} = 0$ as $n \to \infty$ with $\beta_n \to 0$.

Similarly, we can write for $n > 3c$

$$\begin{aligned} \|\mathcal{D}^{\beta_n}_{n,c}(t^2,x) - x^2\|_{\rho_0} &= \sup_{x\in[0,\infty)} \frac{|\mathcal{D}^{\beta_n}_{n,c}(t^2,x) - x^2|}{1+x^2} \\ &\leq \left|\frac{n^2}{(n-2c)(n-3c)(1-\beta_n)^2} - 1\right| \sup_{x\in[0,\infty)} \frac{x^2}{1+x^2} \\ &\quad + \left|\frac{n\left(2-2\beta_n+\beta_n^2\right)}{(n-2c)(n-3c)(1-\beta_n)^3}\right| \sup_{x\in[0,\infty)} \frac{x}{1+x^2} \end{aligned}$$

which implies that

$$\lim_{n\to\infty}\|\mathcal{D}_{n,c}^{\beta_n}(t^2,x)-x^2\|_{\rho_0}=0 \text{ with } \beta_n\to 0.$$

Thus the proof is completed. □

We give the following theorem to approximate all function in $C_{\rho_0}(\mathbb{R}^+)$. This type of results are given in [5] for locally integrable functions.

Theorem 4. *Let $\beta_n\to 0$ as $n\to\infty$. For each $f\in C_{\rho_0}(\mathbb{R}^+)$ and $\lambda>0$, we have*

$$\lim_{n\to\infty}\sup_{x\in[0,\infty)}\frac{|\mathcal{D}_{n,c}^{\beta_n}(f,x)-f(x)|}{(1+x^2)^{1+\lambda}}=0.$$

Proof. For any fixed $x_0>0$,

$$\begin{aligned}\sup_{x\in[0,\infty)}\frac{|\mathcal{D}_{n,c}^{\beta_n}(f,x)-f(x)|}{(1+x^2)^{1+\lambda}} &\le \sup_{x\le x_0}\frac{|\mathcal{D}_{n,c}^{\beta_n}(f,x)-f(x)|}{(1+x^2)^{1+\lambda}}+\sup_{x\ge x_0}\frac{|\mathcal{D}_{n,c}^{\beta_n}(f,x)-f(x)|}{(1+x^2)^{1+\lambda}}\\ &\le \|\mathcal{D}_{n,c}^{\beta_n}(f)-f\|_{C[0,x_0]}\\ &\quad+\|f\|_{\rho_0}\sup_{x\ge x_0}\frac{|\mathcal{D}_{n,c}^{\beta_n}(1+t^2,x)|}{(1+x^2)^{1+\lambda}}+\sup_{x\ge x_0}\frac{|f(x)|}{(1+x^2)^{1+\lambda}}.\end{aligned}$$

The first term of the above inequality tends to zero from Theorem 2. By Lemma 2 for any fixed $x_0>0$ it is easily seen that $\sup_{x\ge x_0}\dfrac{|\mathcal{D}_{n,c}^{\beta_n}(1+t^2,x)|}{(1+x^2)^{1+\lambda}}\to 0$ as $n\to\infty$ with $\beta_n\to 0$. We can choose $x_0>0$ so large that the last part of the above inequality can be made small enough.

Thus the proof is completed. □

Theorem 5. *Let $b>0$ and $\beta_n\in(0,1)$ such that $n\beta_n\to l\in\mathbb{R}$ as $n\to\infty$.*

Then for every $f\in C[0,b]$, twice differentiable at a fixed point $x\in(0,b)$, we have

$$\lim_{n\to\infty}n(D_{n,c}^{\beta_n}(f,x)-f(x))=x(l+2c)f'(x)+\frac{x(2+xc)}{2}f''(x).$$

.

Proof. From Lemma 3 we have

$$\lim_{n\to\infty}n\mu_{1,n,c}^{\beta_n}=x(l+2c),$$

$$\lim_{n\to\infty}n\mu_{2,n,c}^{\beta_n}=x(2+xc),$$

$$\lim_{n\to\infty}n\mu_{4,n,c}^{\beta_n}=0.$$

Using Theorem 2.1 from [1] we get the conclusion. □

4. King's Approach

To make the convergence faster in 2003, King [8] proposed an approach to modify the classical Bernstein polynomial, so that the sequence preserve test functions e_0 and e_2. After this approach many researcher contributed in this direction. Motivated by such type of operators, we generalized King-type approach of the operators (1.3) as follows:

For $\beta \in [0,1)$, $x \in [0,\infty)$ and $f \in C_{\rho_0}(\mathbb{R}^+)$, we introduced King-type positive linear operators as

$$\mathcal{D}^{*\beta}_{n,c}(f,x) \;=\; \frac{(n-c)}{c}\sum_{v=1}^{\infty}\omega_\beta(v,nr_n(x))\int_0^\infty p_{n,v-1}(t)f(t)dt + e^{-nr_n(x)}f(0), \tag{4.1}$$

where $r_n(x) = \dfrac{(n-2c)(1-\beta)x}{n}$ and $\omega_\beta(v,nx)$ and $p_{n,v-1}(t)$ as same as defined in (1.2) and (1.3).

Lemma 4. *The following holds*

$$\mathcal{D}^{*\beta}_{n,c}(1,x) = 1, \qquad \mathcal{D}^{*\beta}_{n,c}(t,x) = x,$$

$$\mathcal{D}^{*\beta}_{n,c}(t^2,x) = \frac{n-2c}{(n-3c)}x^2 + \frac{(2-2\beta+\beta^2)}{(n-3c)(1-\beta)^2}x, \;\; n > 3c,$$

$$\mathcal{D}^{*\beta}_{n,c}(t^3,x) = \frac{x^2n((1-\beta^2)(n-4c)x + 3(\beta^2-2\beta+2))}{(1-\beta)^2(n-3c)(n-4c)} + o\left(\frac{1}{n}\right), \;\; n > 4c,$$

$$\mathcal{D}^{*\beta}_{n,c}(t^4,x) = \frac{n^2x^3((1-\beta)^2(n-6c)x + 6(\beta^2-2\beta+2))}{(1-\beta)^2(n-3c)(n-4c)(n-5c)} + o\left(\frac{1}{n}\right), \;\; n > 5c.$$

The proof follows from Lemma 2 and linearity of $\mathcal{D}^{*\beta}_{n,c}$. We observe that the operators $\mathcal{D}^{*\beta}_{n,c}$ are preserves linear functions that means, $\mathcal{D}^{*\beta}_{n,c}(a+bt,x) = a+bx$ for constants a, b.

Lemma 5. *The following holds*

$$\mu^{*\beta}_{1,n,c} = \mathcal{D}^{*\beta}_{n,c}(t-x,x) = 0,$$

$$\mu^{*\beta}_{2,n,c} = \mathcal{D}^{*\beta}_{n,c}((t-x)^2,x) = \frac{c}{n-3c}x^2 + \frac{(2-2\beta+\beta^2)}{(n-3c)(1-\beta)^2}x, \;\; n > 3c,$$

$$\mu^{*\beta}_{4,n,c} = \mathcal{D}^{*\beta}_{n,c}((t-x)^4,x) = o\left(\frac{1}{n}\right), \;\; n > 5c.$$

The proof follows from Lemma 4.

Theorem 6. *Let $\mathcal{D}^{*\beta_n}_{n,c}, n > 3c$, be defined as in* (4.1)*, where $\lim_{n\to\infty}\beta_n = 0$. For any compact $B \subset \mathbb{R}^+$ and for each $f \in C_{\rho_0}(\mathbb{R}^+)$ one has*

$$\lim_{n\to\infty}\mathcal{D}^{*\beta_n}_{n,c}(f,x) = f(x), \;\; \textit{uniformly in } x \in B.$$

The proof follows from Lemma 4 and Korovkin theorem [9].

Theorem 7. *For $f \in C_B[0,\infty)$, $0 < \beta < 1$ and $n > 3$, we have*

$$|\mathcal{D}^{*\beta}_{n,c}(f,x) - f(x)| \le C_1\omega_2\left(f, \sqrt{\hat{\mu}^{\beta}_{n,2,c}}\right),$$

where C_1 is a positive constant.

Proof. Let $g \in W^2_\infty$ and $x \in [0,\infty)$. Using Taylor's expansion

$$g(t) = g(x) + g'(x)(t-x) + \int_x^t (t-u)g''(u)du, \quad t \in [0,\infty)$$

and Lemma 4, we have

$$\mathcal{D}_{n,c}^{*\beta}(g,x)-g(x)=\mathcal{D}_{n}^{*\beta}\left(\int_x^t (t-u)g''(u)du\right).$$

Notice that, $\left|\int_x^t (t-u)g''(u)du\right| \leq (t-x)^2\|g''\|$. Thus

$$|\mathcal{D}_{n,c}^{*\beta}(g,x)-g(x)| \leq \mathcal{D}_{n,c}^{*\beta}((t-x)^2,x)\|g''(x)\| = \mu_{2,n,c}^{*\beta}\|g''\|.$$

Since $|\mathcal{D}_{n,c}^{*\beta}(f,x)| \leq \|f\|$,

$$\begin{aligned}|\mathcal{D}_{n,c}^{*\beta}(f,x)-f(x)| &\leq |\mathcal{D}_{n,c}^{*\beta}(f-g,x)-(f-g)(x)|+|\mathcal{D}_{n,c}^{*\beta}(g,x)-g(x)| \\ &\leq 2\|f-g\|+\mu_{2,n,c}^{*\beta}\|g''\|.\end{aligned}$$

Finally taking the infimum on right side over all $g \in W_\infty^2$ and using (3.2), we get the desired result. ☐

Now, we establish the Voronovskaja type asymptotic formula for the operators $\mathcal{D}_{n,c}^{*\beta}$:

Theorem 8. *Let $b > 0$ and $\beta_n \in (0,1)$ such that $n\beta_n \to 0$ as $\to \infty$.*

Then for every $f \in C[0,b]$, twice differentiable at a fixed point $x \in (0,b)$, we have

$$\lim_{n\to\infty} n\big(\mathcal{D}_{n,c}^{*\beta_n}(f,x)-f(x)\big) = x(2+xc)\frac{f''(x)}{2}.$$

Proof. From Lemma 5 we have

$$\lim_{n\to\infty} n\mu_{1,n,c}^{*\beta_n}=0,$$

$$\lim_{n\to\infty} n\mu_{2,n,c}^{*\beta_n}=x(2+xc),$$

$$\lim_{n\to\infty} n\mu_{4,n,c}^{*\beta_n}=0.$$

Using Theorem 2.1 from [1] we get the conclusion. ☐

References:

References

[1] D. Cardenas-Morales, P. Garrancho, I. Rasa, *Asymptotic formulae via a Korovkin-type result*, Abstract and Applied Analysis, Volume 2012, Article ID 217464, 12 pages.

[2] D.K. Dubey, V.K. Jain, *Rate of approximation for integrated Száasz-Mirakyan operators*, Demonstr. Math. **41:4** (2008), 879–886.

[3] A. Farcas, *An asymptotic formula for Jain's operators*, Stud. Univ. Babes Bolyai Math., **57** (2012), No. 4, 511–517.

[4] A.D. Gadjiev, *Theorems of the type of PP Korovkin theorems*, Mat. Zametki **20:5** (1976), 781–786.

[5] A.D. Gadjiev, R.O. Efendiyev, E. Ibikli, *On Korovkin type theorem in the space of locally integrable functions*, Czechoslov. Math. J. **53:1** (2003), 45–53.

[6] V. Gupta, Themistocles M. Rassias, *Direct estimates for certain Szász type operators*, Applied Mathematics and Computation **251** (2015), 469–474.

[7] G.C. Jain, *Approximation of functions by a new class of linear operators* J. Aust. Math. Soc. **13:3**, (1972) 271–276.

[8] J.P. King, *Positive linear operators which preserve* x^2, Acta Math. Hung. **99:3** (2003), 203–208.

[9] P.P. Korovkin, *On convergence of linear positive operators in the space of continuous functions*, Dokl. Akad. Nauk SSSR **90** (1953), 961–964.

[10] P. Patel, V. N. Mishra, *Jain-Baskakov Operators and its Different Generalization*, Acta Math Vietnam DOI 10.1007/s40306-014-0077-9.

[11] R.A. DeVore, G.G. Lorentz, *Constructive Approximation. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*, 303. Springer, Berlin. x+449 pp (1993).

[12] O. Szász, *Generalization of S. Bernstein polynomials to the infinite interval*, J. Res. Natl. Bur. Stand. **45** (1950) 239–245.

[13] S. Tarabie, *On Jain-Beta linear operators*, Appl. Math. Inf. Sci. **6:2**, (2012) 213–216.

[14] S. Umar, Q. Razi, *Approxiamtion of function by generalized Szász operators*, Communications de la Faculté Des Sciences de l'Université Ankara Mathematique, **34** (1985), 45–52.

[15] H. Gonska, P. Piţul, I. Raşa, *General King-Type Operators*, Result. Math. **53**, Issue 3-4 (2009), 279–286.

[16] H. Gonska, D. Kacsó, I. Raşa, *On Genuine Bernstein-Durrmeyer Operators*, Result. Math. **50** (2007), 213–225.

[17] H. Gonska, I. Raşa, *Asymptotic behaviour of differntiated bernsein polynomials*, matematički vesnik, **61** (2009), 53-60.

[18] T. Acar, A. Aral, I. Raşa, *Modified Bernstein Durrmeyer operators*, General Mathematics, **22** (1), 2014, 27-41.